

Isotope studies in rock phosphates

Miyuki Maekawa¹, Roland Bol², Yajie Sun², Liankai Zhang¹, Silvia Haneklaus¹ and Ewald Schnug¹

¹*Institute for Crop and Soil Science, Julius Kühn-Institut, Federal Research Institute for Cultivated Plants, Bundesallee 50, Braunschweig, Germany (E-mail: miyuki.maekawa@julius-kuehn.de);* ²*Forschungszentrum Jülich IBG-3, Wilhelm-Johnen-Straße, 52428 Jülich, Germany*

Isotope studies comprise tracer experiments and the determination of isotope ratios in fertilizers. So, it is for instance possible to discriminate crops in relation to soil characteristics and fertilizer type which have been labeled with the ³⁴S isotope by calculating $\delta^{34}\text{S}$ after measuring the ratio of stable ³⁴S/³²S isotope masses. The determination of the ⁸⁷Sr/⁸⁶Sr ratio proved to be suitable to distinguish phosphates bound in phosphorites from those in carbonatites. Such discrimination delivers further information about the contamination with heavy metals or rare earth elements which can be beneficiated and taken as raw material for industrial purposes. A major obstacle in fertilizer research is that the effective, long-term utilization of phosphate cannot be determined empirically as P is an isotopic with only one stable isotope, ³¹P so that fractionation studies are not feasible. Radioactive P tracers cannot be employed either as the half-life time of ³³P is with only 25.3 days and that of ³²P 14.3 days too short for long-term experiments. Accordingly, the geological origin of rock phosphates cannot be determined by stable P isotope ratios. Here, the $\delta^{18}\text{O}$ and activity ratio of ²³⁴U/²³⁸U are used to monitor provenances. It is the objective of this study to outline the current status of isotopic studies in rock phosphates, to summarize the significance of these data and to depict future analytical options in order to enable a proper attribution of the origin of rock phosphates and to follow up fluxes of contaminants in the environment.

Bearbeitet von/ Compiled by:

Luit J. De Kok², Silvia Haneklaus¹, Ewald Schnug¹

**25th International Symposium of the Scientific
Centre for Fertilizers “Significance of Sulfur in
High-Input Cropping Systems”
Groningen (Netherlands), September 5-8, 2017**

¹Julius Kühn-Institut (JKI), Bundesforschungsinstitut für Kulturpflanzen
Institut für Pflanzenbau und Bodenkunde

²University of Groningen
Faculty of Science and Engineering

Berichte aus dem Julius Kühn-Institut

191



Kontaktadresse/ Contact

Dr. Silvia Haneklaus
Federal Research Centre for Cultivated Plants
Institute for Crop and Soil Science
Bundesallee 50
38116 Braunschweig
Germany

Telefon +49 (0) 531 596 2121
Telefax +49 (0) 531 596 2199

Wir unterstützen den offenen Zugang zu wissenschaftlichem Wissen.
Die Berichte aus dem Julius Kühn-Institut erscheinen daher als OPEN ACCESS-Zeitschrift.
Alle Ausgaben stehen kostenfrei im Internet zur Verfügung:
<http://www.julius-kuehn.de> Bereich Veröffentlichungen – Berichte.

We advocate open access to scientific knowledge. Reports from the Julius Kühn Institute are therefore published as open access journal. All issues are available free of charge under <http://www.julius-kuehn.de> (see Publications – Reports).

Herausgeber / Editor

Julius Kühn-Institut, Bundesforschungsinstitut für Kulturpflanzen, Braunschweig, Deutschland
Julius Kühn Institute, Federal Research Centre for Cultivated Plants, Braunschweig, Germany

Vertrieb

Saphir Verlag, Gutsstraße 15, 38551 Ribbesbüttel
Telefon +49 (0)5374 6576
Telefax +49 (0)5374 6577

ISSN 1866-590X

DOI 10.5073/berjki.2017.191.000



Dieses Werk ist lizenziert unter einer [Creative Commons – Namensnennung – Weitergabe unter gleichen Bedingungen – 4.0 Lizenz](https://creativecommons.org/licenses/by-sa/4.0/).
This work is licensed under a [Creative Commons – Attribution – ShareAlike – 4.0 license](https://creativecommons.org/licenses/by-sa/4.0/).